



## Full Length Research Article

Advancements in Life Sciences – International Quarterly Journal of Biological Sciences

## ARTICLE INFO

## Open Access



Date Received:  
08/02/2021;  
Date Revised:  
15/11/2024;  
Available Online:  
31/12/2024;

# *In vitro* step up NaCl treatment method as inducer of salinity tolerance in Basmati rice varieties

## Author's Affiliation:

Institute of Botany, University of  
the Punjab, Lahore – Pakistan

Naveeda Batool\*, Humera Afrasiab

## \*Corresponding Author:

Naveeda Batool  
Email:  
[hellonaveeda@yahoo.com](mailto:hellonaveeda@yahoo.com)

## How to Cite:

Batool N, Afrasiab H (2025).  
*In vitro* step up NaCl  
treatment method as inducer  
of salinity tolerance in  
Basmati rice varieties. Adv.  
Life Sci. 12(1): 168-174.

## Keywords:

*In vitro* selection; Salt  
tolerance; Basmati rice; Step  
up NaCl treatment;  
Callogenesis; Plant  
regeneration

## Editorial Note:

You are viewing the latest  
version of this article having  
minor corrections related to  
the use of English language.

## Abstract

**Background:** Salt has been recorded to negatively impacts rice crop's seedling and reproductive stages. Globally, soil salinity is 2<sup>nd</sup> most important abiotic factor, profoundly impacting rice productivity. The purpose of current research is to induce salinity tolerance in salt sensitive varieties of Basmati rice [Basmati-385 (Bas-385), Super Basmati (Sup-Bas) and Basmati-2000 (Bas-2000)] by use of *in vitro* step-up NaCl treatment method.

**Methods:** The *in vitro* step up NaCl treatment method was used to produce salt-tolerant calli from salt-sensitive Basmati rice varieties. The regeneration of these salt-tolerant calli in a salt-free medium led to the M<sub>1</sub> generation. Subsequently, M<sub>2</sub> and M<sub>3</sub> generations were cultivated and grown under salt stress conditions.

**Results:** The highest callus induction percentage was achieved for all Basmati rice varieties by using MS medium supplemented with 2.0 mg/L 2,4-D. These calli underwent *in vitro* step up NaCl treatments, with the maximum tolerance level to NaCl being 1.0% across all varieties. Bas-385 and Bas-2000 exhibited regeneration frequencies of 75% and 85% respectively when cultured in MS medium supplemented with 2.0 mg/L BAP and 0.5 mg/L NAA, while Sup-Bas displayed a regeneration frequency of 79% in MS medium comprising 3.0 mg/L BAP with 0.5 mg/L NAA. In subsequent M<sub>2</sub> and M<sub>3</sub> generations, a decline was observed in all selected agronomic and morphological features in all varieties under salinity stress in comparison to parent plants.

**Conclusion:** This research validates the induction of salt tolerance in salt susceptible varieties of Basmati rice by *in vitro* step up NaCl treatment technique.



## Introduction

Rice (*Oryza sativa* L.) is cultivated extensively worldwide, serving as a staple food for over half of the global population and contributing to 21% of global per capita energy consumption [1]. It is rich in dietary energy (27%), protein (20%), and fat (3%) [2]. Rice is grown across all continents except Antarctica, covering more than 167.13 million hectares and yielding approximately 782 million tons [3,4]. However, the majority of rice production occurs in Asia, accounting for over 90.7% of the world's output [5,3]. In Pakistan, rice cultivation occupies about 11% (3,034 thousand hectares) of the total cultivated area, yielding 7,410 thousand tons annually. It is a high-value cash crop, contributing 0.6% to value added in the gross domestic product and 3.1% to the agricultural sector [6].

The Basmati aromatic traditional rice and the short-stemmed and high-yielding IRRI-Pak varieties are the two main types of rice grown in Pakistan. The most prevalent type, aromatic Basmati rice, is found in 62% of the country's rice fields and accounts for 52% of the country's rice production. Basmati varieties are primarily cultivated in the Punjab province, where 94% of the Basmati area and 92% of the production are concentrated [6]. In Punjab, there has been a significant increase of 60% in Basmati cultivation area and a staggering 155% increase in yield from 1982 to 2002 [6,7].

Despite its genetic potential, rice yield has not reached its maximum in most rice-growing regions globally. One of the key reasons for this shortfall is the susceptibility of rice to various biotic (pests, insects, diseases) and abiotic factors (soil salinity, drought, waterlogging, cold) [8,9]. Soil salinity ranks as the second most crucial abiotic factor affecting rice productivity worldwide [10], with rice being recognized as salt-susceptible, leading to adverse effects during both sprouting and reproductive stages [11-15].

Salt-affected soils pose a significant challenge at universal and local farm levels. Out of the total 167.13 million hectares of rice-growing land, approximately 31% is afflicted with high salinity levels, greatly impeding normal rice productivity. In Pakistan, one million hectares of rice-growing region is pretentious by salinity, consequentially lead to a decline of 40–70% in crop yield [16-19].

Recent projections estimate the global population to reach 9 billion by 2050, with emerging countries contributing 97% of this growth. Rice demand is expected to reach 590 million tons by 2050 [20,2]. To meet this escalating demand, developing new rice varieties capable of growing under salt stress by exploiting genetic variability is imperative.

The use of plant tissue culture technology for screening plants tolerant to abiotic stresses offers a

promising approach for selecting cell populations on a large scale [21]. *In vitro* screening methods allow rapid proliferation of cell populations in petri dishes or flasks using well-defined growth media. Cell lines established through tissue culture applications can be vetted *in vitro* for tolerance to various stresses, including salinity, which has been widely employed in rice breeding [22-24] and other plants [25,21].

Soil salinity is now recognized as a significant problem in Pakistan's prime Basmati rice-growing areas [26], highlighting the need for developing rice varieties resistant to salinity effects. The purpose of this study is to use the *in vitro* step-up NaCl treatment method to persuade the salinity tolerance in economically imperative salt susceptible varieties of Basmati rice (Bas-385, Sup-Bas and Bas-2000).

## Methods

### Explants disinfection

Licensed seeds of Bas-385, Sup-Bas, and Bas-2000 underwent a cleaning process by washing with cleanser for 15 minutes, followed by several rinses with tap water. Subsequently, the seeds were drenched in distilled water overnight. For surface sterilization, a 0.1% HgCl<sub>2</sub> (Mercuric chloride) solution was employed, and manually dehusked seeds were soaked for 15 minutes. Afterwards, the seeds underwent a thorough washing with sterilized purified water, repeated 3-5 times within the sterilized atmosphere of a laminar airflow cabinet.

### Callogenesis

For callus induction, disinfected seeds were introduced onto agar-solidified MS [27] medium supplemented with various combinations of plant growth regulators [2,4-Dichlorophenoxy acetic acid (2,4-D), Benzyl aminopurine (BAP), Kinetin (KIN)], with pH attuned to 5.7 - 5.8. The cultures were incubated in darkness at 25 ± 1°C temperature. Callus cultures were allowed to develop for 4 weeks. at the end of the 4th week, information regarding frequency of callus induction and the morphology of callus were recorded. The medium exhibiting the highest prospective for callus formation and proliferation rate was identified as the optimal medium for next experimentation. Non-embryogenic calli were discarded, while vigorous embryogenic calli were carefully chosen and sub-cultured onto new medium for proliferation after every 2 weeks.

### *In vitro* selection of salt tolerance calli

Vigorous embryogenic calli, aged eight weeks, were carefully excised under aseptic conditions and exposed to *in vitro* step-up NaCl treatments method. These treatments involved a range of NaCl concentrations

from 0, 0.5, 1, 1.5, 2 to 2.5%, incorporated into agar coagulated Murashige and Skoog medium, at intermissions of 3 weeks. The calli exhibiting high salt tolerance were maintained on the same medium for three sequential proliferation phases, each phase lasting for six weeks.

### Plant regeneration

To initiate regeneration, all NaCl-tolerant calli were transferred to MS medium supplemented with diverse recipes of growth regulators. The cultures were then cultivated in a growth room under a photoperiod of 16 hrs. light and 8 hrs. dark at  $25\pm1^{\circ}\text{C}$  temperature. After every three weeks, the cultures were shifted to new fresh medium. The appearance of green shoots measuring over 3cm in height was considered for calculating the Regeneration Frequency (RF).

These regenerated calli were transferred to basic Murashige and Skoog medium devoid of any growth regulators to promote root development. Subsequently, they were shifted to a hydroponic solution to facilitate root hardening. Each regenerated plant was allotted a unique plant line number ( $M_1$ ) and allowed to grow until reaching maturity.

### Screening for salt tolerance from $M_2$ generation

$M_2$  lines derived from  $M_1$  plants, excluding those exhibiting poor plant characteristics such as droopy leaves and weak culms, were harvested and sequentially numbered from  $M_2$  1 to n. Screening for salt tolerance was conducted by cultivating all  $M_2$  lines under saline field conditions, with an electrical conductivity (EC) equivalent to the maximum percentage of NaCl (14 ds/m) that the salt-tolerant calli had previously survived.

### Screening for salt tolerance from $M_3$ generation

After 140 days, the  $M_2$  generation was harvested, now referred to as the  $M_3$  generation, exhibiting normal grain fertility at a rate of 80% on an individual plant basis.  $M_3$  lines were sequentially numbered from  $M_3$  1 to n. Screening for salinity tolerance in the  $M_3$  lines was conducted by cultivating all lines in a salt stressed field with an electrical conductivity (EC) equal to the maximum percentage of NaCl (14 ds/m) that the salt-tolerant calli had previously endured. Control plants (parent plants) of all varieties were also transplanted alongside the  $M_3$  lines for comparison.

### Performance of $M_2$ and $M_3$ generations

Data regarding several agronomic and morphological traits of the salt-tolerant  $M_2$  and  $M_3$  lines were meticulously recorded for all varieties. The selected traits included height of plant, length of panicle, number of tillers and the number of plants producing fertile seeds.

### Statistical analysis


All experimentation was conducted in a triplicate manner. The obtained data was subjected to Analysis of Variance (ANOVA), and the differences between the means were deliberate through Duncan's Multiple Range Test at a significance level of  $P < 0.05$ . SPSS version 12 was used to do the statistical analysis.

## Results

### Callogenesis

Callogenic response of Basmati rice varieties on Murashige and Skoog (MS) medium supplemented with various combinations of plant growth regulators after 4 weeks of inoculation is given in Table 1. No callogenic response was observed in control and the maximum callus induction response in Basmati-385 (78%), Super Basmati (75%) and Basmati-2000 (80%) was achieved on MS medium fortified with 2.0 mg/L 2,4-D. The calli were granular and compact in texture, embryogenic in nature and light yellow to light brown in color (Fig. 1a, 1b and 1c). The second-best results were obtained in all the varieties in Murashige and Skoog medium containing 1.0 mg/L 2,4-Dichlorophenoxy acetic acid while the percentage of callus induction decreased across all varieties as the concentration of 2,4-D increased.

When combining 2,4-D (2.0 mg/L) with different levels of BAP, the highest callus induction was observed on BAP at a level of 2.0 mg/L across all varieties. However, this induction decreased with further increases in BAP concentration. Similarly, when different concentrations of KIN were tested with 2,4-Dichlorophenoxy acetic acid (2.0 mg/L), the optimal retort was obtained at a KIN concentration of 1.0 mg/L in all varieties (refer to Table 1).



**Figure 1:** Callogenic response of (a) Basmati-385, (b) Super Basmati and (c) Basmati-2000 on Murashige and Skoog medium invigorated with 2.0 mg/L 2,4-Dichlorophenoxy acetic acid.

### In vitro selection of salt tolerance calli

Vigorous embryogenic calli of Bas-385, Sup-Bas and Bas-2000, aged eight weeks, were aseptically cut out. Approximately 50mg (initial weight) of calli from every variety were exposed to *in vitro* step-up NaCl treatments. These treatments ranged from 0, 0.5, 1, 1.5, 2 to 2.5% NaCl, incorporated into agar-solidified Murashige and Skoog medium invigorated with 2.0 mg/L 2,4-Dichlorophenoxy acetic acid. The treatments were administered at 3-week intervals (Table 2).

Relative growth value of calli in each variety was decreased with increasing NaCl concentration as compared to control. In case of Bas-385, relative growth at 0.5% NaCl was 1.90 and 1.95 at 1.0% NaCl. While, at higher concentration of NaCl (1.5, 2.0 and 2.5%) growth value sharply decreased, indicating that maximum NaCl tolerance level for calli of Bas-385 was 1.0%. Similar trend for percentage NaCl tolerance level was also observed in case of Sup-Bas and Bas-2000 (Table 2).

| Medium                     | Plant Growth regulators (mg/L) | Callus induction(%)         |                             |                             |
|----------------------------|--------------------------------|-----------------------------|-----------------------------|-----------------------------|
|                            |                                | Bas-385                     | Sup-Bas                     | Bas-2000                    |
| MS + 2,4-D                 | 0                              | 00                          | 00                          | 00                          |
|                            | 1                              | 66.33 ± 4.25 <sup>b</sup>   | 60.45 ± 6.66 <sup>bc</sup>  | 70.65 ± 3.41 <sup>b</sup>   |
|                            | 2                              | 78.41 ± 6.23 <sup>a</sup>   | 75.77 ± 6.45 <sup>a</sup>   | 80.88 ± 7.80 <sup>a</sup>   |
|                            | 3                              | 43.26 ± 2.12 <sup>cd</sup>  | 45.23 ± 4.25 <sup>cd</sup>  | 50.45 ± 4.25 <sup>bc</sup>  |
|                            | 4                              | 35.32 ± 3.33 <sup>d</sup>   | 40.42 ± 4.10 <sup>cd</sup>  | 30.33 ± 3.25 <sup>de</sup>  |
|                            | 5                              | 25.55 ± 2.12 <sup>def</sup> | 25.22 ± 2.36 <sup>def</sup> | 20.14 ± 3.41 <sup>efg</sup> |
|                            | 6                              | 15.11 ± 2.41 <sup>ef</sup>  | 15.11 ± 1.25 <sup>ef</sup>  | 10.11 ± 1.24 <sup>fg</sup>  |
|                            | 7                              | 10.12 ± 1.14 <sup>ef</sup>  | 10.12 ± 1.22 <sup>ef</sup>  | 10.11 ± 1.24 <sup>fg</sup>  |
| MS + 2,4-D (2.0mg/L) + BAP | 8                              | 08.25 ± 1.36 <sup>efg</sup> | 09.11 ± 1.00 <sup>efg</sup> | 10.11 ± 1.24 <sup>fg</sup>  |
|                            | 1                              | 41.14 ± 2.45 <sup>b</sup>   | 36.25 ± 3.36 <sup>c</sup>   | 49.44 ± 4.25 <sup>b</sup>   |
|                            | 2                              | 45.45 ± 2.36 <sup>a</sup>   | 40.14 ± 4.12 <sup>bc</sup>  | 53.32 ± 3.25 <sup>a</sup>   |
|                            | 3                              | 32.11 ± 2.05 <sup>bc</sup>  | 31.18 ± 3.41 <sup>cd</sup>  | 40.41 ± 4.12 <sup>c</sup>   |
| MS + 2,4-D (2.0mg/L) + KIN | 4                              | 20.22 ± 4.12 <sup>de</sup>  | 23.41 ± 2.11 <sup>def</sup> | 30.22 ± 3.21 <sup>de</sup>  |
|                            | 1                              | 43.41 ± 3.22 <sup>ab</sup>  | 53.23 ± 5.41 <sup>a</sup>   | 48.41 ± 4.26 <sup>b</sup>   |
|                            | 2                              | 31.33 ± 2.45 <sup>bc</sup>  | 36.25 ± 3.23 <sup>c</sup>   | 33.10 ± 3.21 <sup>de</sup>  |
|                            | 3                              | 28.41 ± 1.45 <sup>cd</sup>  | 30.32 ± 1.25 <sup>cd</sup>  | 24.32 ± 2.33 <sup>ef</sup>  |
|                            | 4                              | 25.71 ± 1.32 <sup>cd</sup>  | 26.45 ± 2.25 <sup>de</sup>  | 10.12 ± 3.02 <sup>fg</sup>  |

**Table 1:** Callogenic expression of Basmati rice varieties in MS medium invigorated with different plant growth regulators. 100 seeds of each variety were inoculated in each experiment. Experiments were performed in triplicate.

| Basmati Rice variety | NaCl Treatments (%) | Relative growth (%) |
|----------------------|---------------------|---------------------|
| Bas-385              | Control             | 2.20 ± 1.02a        |
|                      | 0.5 %               | 1.90 ± 0.88b        |
|                      | 1.0%                | 1.95 ± 0.75b        |
|                      | 1.5%                | 0.85 ± 0.21c        |
|                      | 2.0%                | 0.71 ± 0.11cd       |
| Sup-Bas              | 2.5%                | 0.40 ± 0.09d        |
|                      | Control             | 2.40 ± 0.98a        |
|                      | 0.5 %               | 1.73 ± 0.66b        |
|                      | 1.0%                | 1.80 ± 0.85b        |
|                      | 1.5%                | 0.73 ± 0.23cd       |
| Bas-2000             | 2.0%                | 0.55 ± 0.41cd       |
|                      | 2.5%                | 0.29 ± 0.10de       |
|                      | Control             | 2.30 ± 1.00a        |
|                      | 0.5 %               | 1.86 ± 1.04b        |
|                      | 1.0%                | 1.93 ± 1.11b        |
|                      | 1.5%                | 0.84 ± 0.32c        |
|                      | 2.0%                | 0.62 ± 0.11cd       |
|                      | 2.5%                | 0.36 ± 0.09cde      |

Means values trailed by the diverse letters are meaningfully different as per DMRT p<0.05.

**Table 2:** Assortment of NaCl tolerant calli by use of step up NaCl treatment method.

(1) Around 50mg of callus (initial weight) underwent *in vitro* step-up NaCl treatment, spanning concentrations from 0, 0.5, 1, 1.5, 2 to 2.5% NaCl. This treatment was administered in agar coagulated MS medium supplemented with 2 mg/L 2,4-

Dichlorophenoxy acetic acid at three week intermissions.

(2) Relative growth: The ratio of the calli's final weight to its initial weight is what determines their relative growth. The outcomes that have been reported are the average of three replicates.C

### Plant regeneration

The calli of Basmati rice varieties, which exhibited high tolerance to salt (1.0% NaCl), underwent three consecutive proliferation phases in the same Murashige and Skoog medium invigorated with 2,4-Dichlorophenoxy acetic acid 2.0 mg/L and 1.0% NaCl, with a 6-week interval each time, to fully adapt to this level of salinity stress. For regeneration, all chosen salt-tolerant calli were transferred to Murashige and Skoog medium supplemented with different blends of 6-Benzylaminopurine with 0.5mg/L 1-Naphthaleneacetic acid. A hundred calli from each variety were moved to the regeneration medium. According to Table 3, in Murashige and Skoog medium invigorated with 2.0 mg/L 6-Benzylaminopurine and 0.5 mg/L 1-Naphthaleneacetic acid, Bas-385 and Bas-2000 had regeneration frequencies of 75% and 85%, respectively. Sup-Bas had regeneration frequency of 79% in Murashige and Skoog medium invigorated with 3.0 mg/L 6-Benzylaminopurine and 0.5 mg/L 1-Naphthaleneacetic acid. It was noted that the percentage of regeneration frequency decreased as BAP levels rose (Table 3).

| Medium                           | Regeneration frequency (%) |              |              |
|----------------------------------|----------------------------|--------------|--------------|
|                                  | Bas-385                    | Sup-Bas      | Bas-2000     |
| MS + 0.5mg/LNAA + BAP (1.0 mg/L) | 60.65±5.22c                | 58.45±4.36bc | 63.36±6.33c  |
| MS + 0.5mg/LNAA + BAP (2.0 mg/L) | 75.76±4.57a                | 60.62±2.12bc | 85.14±7.65a  |
| MS + 0.5mg/LNAA + BAP (3.0 mg/L) | 70.41±7.22b                | 79.71±4.65a  | 74.23±4.56b  |
| MS + 0.5mg/LNAA + BAP (4.0 mg/L) | 52.31±2.64d                | 41.21±5.32c  | 50.54±5.23d  |
| MS + 0.5mg/LNAA + BAP (5.0 mg/L) | 38.25±4.25ef               | 30.33±2.33cd | 24.21±1.23ef |

Means values trailed by the different letters are meaningfully different according to DMRT p<0.05.

**Table 3:** Rejuvenation potential of 1.0% NaCl Tolerant calli.

The shoots grown *in vitro* were shifted to MS basal medium (barren of any growth regulators) to induce root formation (refer to Table 4). Subsequently, the *in vitro* grown roots were transferred to a hydroponic solution for the purpose of hardening. The plant line number (M<sub>1</sub>) was given to each regenerated plant, and it was allowed to grow until it reached maturity. Later M<sub>2</sub> and M<sub>3</sub> generation were also obtained and the number of plants per variety in each generation is given in Table 4.

| Basmati varieties | Number of plants        |                          |                          |
|-------------------|-------------------------|--------------------------|--------------------------|
|                   | M <sub>1</sub>          | M <sub>2</sub>           | M <sub>3</sub>           |
| Bas-385           | 70.24±4.35 <sup>a</sup> | 50.54±5.54 <sup>a</sup>  | 35.35±2.35 <sup>a</sup>  |
| Sup-Bas           | 72.42±6.13 <sup>b</sup> | 45.63±4.13 <sup>bc</sup> | 20.42±2.24 <sup>b</sup>  |
| Bas-2000          | 78.75±7.53 <sup>a</sup> | 60.65±6.16 <sup>a</sup>  | 32.23±2.35 <sup>ab</sup> |

Means values trailed by the different letters are expressively different according to DMRT  $p < 0.05$ .

**Table 4:** Number of salt tolerant plants attained in M<sub>1</sub>, M<sub>2</sub> and M<sub>3</sub> generations.

| Varieties     | Plant height (cm) |                |                | Panicle length |                |                | No of Tillers |                |                | No of plants with fertile seeds |                |                | Total weight (g) <sup>1</sup> |                |                |
|---------------|-------------------|----------------|----------------|----------------|----------------|----------------|---------------|----------------|----------------|---------------------------------|----------------|----------------|-------------------------------|----------------|----------------|
|               | Parent            | M <sub>2</sub> | M <sub>3</sub> | Parent         | M <sub>2</sub> | M <sub>3</sub> | Parent        | M <sub>2</sub> | M <sub>3</sub> | Parent                          | M <sub>2</sub> | M <sub>3</sub> | Parent                        | M <sub>2</sub> | M <sub>3</sub> |
| Basmati-385   | 133               | 85<br>(36)     | 73<br>(45.1)   | 30             | 24<br>(20.0)   | 20<br>(33.0)   | 25            | 16<br>(36.0)   | 12<br>(52.0)   | 50                              | 35<br>(30.0)   | 21<br>(58)     | 420                           | 290<br>(30.9)  | 184<br>(56.1)  |
| Super Basmati | 115               | 67<br>(41.7)   | 53<br>(53.9)   | 25             | 19<br>(24.0)   | 16<br>(36.0)   | 20            | 13<br>(35.0)   | 8<br>(60.0)    | 50                              | 20<br>(60)     | 10<br>(80)     | 432                           | 163<br>(62.2)  | 89<br>(79.3)   |
| Basmati-2000  | 134               | 75<br>(44)     | 67<br>(50)     | 32             | 25<br>(21.8)   | 22<br>(31.2)   | 27            | 18<br>(33.3)   | 14<br>(48.1)   | 50                              | 32<br>(36)     | 19<br>(62)     | 450                           | 270<br>(40.0)  | 168<br>(62.6)  |

<sup>1</sup>Total weight was measured with all plants with fertile seeds.

**Table 5:** Depiction of salt tolerant M<sub>2</sub> and M<sub>3</sub> generations.

### Performance of M<sub>2</sub> and M<sub>3</sub> generations

Morphological and agronomical characteristics of M<sub>2</sub> and M<sub>3</sub> generations were documented and compared with those of parent plants. A decline in all features was noted in all varieties under saline conditions in comparison to the parent plants (refer to Table 5). Results indicate that Sup-Bas exhibited the highest sensitivity to salt stress, displaying a significant decrease in nearly all morphological and agronomical traits considered in the M<sub>2</sub> and M<sub>3</sub> generations. Among the rest of 2 varieties, Bas-385 demonstrated greater salt resistance and exhibited superior field performance compared to Bas-2000.

## Discussion

Abiotic stresses are the primary constraint to development and yield of plants in the farming area. Among them, salt stress is a main pressing issue in arid and semi-arid regions of the world. The detrimental effects that salinity has on plant development include osmotic and oxidative stress, mineral deficiency, nutritional imbalance, or a combination of these factors [28, 29].

Tissue culture technique is being routinely employed for screening of stress tolerance cell lines in various plant species. Using salts as a selective agent, the *in vitro* selection pressure method has been successfully used to induce tolerance to salinity in plants, allowing the preferred survival and growth of desired genotypes [30].

In the present work, from mature seed explants, embryogenic calli of three varieties of Basmati rice (Bas-385, Sup-Bas, and Bas-2000) were induced. The MS medium supplemented with 2.0 mg/L 2,4-D was found to be the best medium showing maximum callus induction and proliferation potential for all these Basmati rice varieties as compared to MS medium containing 2,4-D in mix with various concentrations of BAP or KIN. 2,4-Dichlorophenoxy acetic acid at the level of 2.0 mg/L has also been reported by Ullah *et al.*

inducer growth regulator for Basmati-385 followed by Basmati-370 [31]. Similar findings were reported by Afrasiab and Jafar working with Super Basmati and Rafique *et al.* working with Basmati-370 [32,33].

Several researchers have also reported that 2,4-D at 2.0 mg/L produced the most desired calli in different varieties of rice [34 - 38]. Therefore, it can be concluded that that 2,4-D alone is more effective for the induction of callus in rice varieties as compared to combination of hormones in different concentrations. Upadhyaya *et al.* also reported 2,4-D as the only growth regulator effective in callus induction media for rice [39].

In this research work, the *in vitro* step up NaCl treatment method resulted in calli of Basmati rice varieties that were 1% NaCl-tolerant. While at higher concentration of NaCl (1.5, 2.0 and 2.5%) the calli turned brown and eventually died. Same method was also used by Miki *et al.* for induction of salinity tolerance in salt sensitive rice cultivar Nipponbare and vigorously growing shoot bud clumps up to 2% NaCl tolerant were obtained [22]. Our results are well supported by Htwe *et al.* who described that increasing the concentration of NaCl into culture medium resulted in necrosis of callus in five rice genotypes [40]. Likewise, Wu *et al.* explained that the browning of callus may be due to necrosis or tissue damage as result of oxidation of phenolic compounds [41]. *In vitro* step up NaCl treatment method is an alternative approach for induction of salinity tolerance in calli or plantlets as compared to single step salt treatment [22].

The salt tolerant calli were transferred to regeneration medium and best response was found to be 85% for Basmati-2000 and 75% for Basmati-385 in MS medium containing NAA (0.5 mg/l) with BAP (2.0 mg/l) and 79% for Super Basmati in MS medium having NAA(0.5 mg/l) with BAP (3.0 mg/l). Similar combination of hormones have been reported by Singh *et al.* for plant regeneration of four varieties of rice

from callus [34]. Regenerates of these salt tolerant calli were grown as M<sub>1</sub> generation in salt free field conditions. M<sub>2</sub> and M<sub>3</sub> generation were grown in saline field with EC 14 dS/m which was equal to maximum % of salt which was tolerated by the calli of different rice varieties.

All the agronomical and morphological characters studied showed a significant decrease as compared to control plants. Reduced growth parameters due to salt stress are a common occurrence in plants, observed in cultured cells, tissues, or organs grown on NaCl-supplemented medium. Slower growth is often considered an adaptive mechanism for plant survival under stress, with salt tolerance being inversely related to growth rate [42]. Khorami and Safarnejad found that increased NaCl levels elevated Na<sup>+</sup> and Cl<sup>-</sup> concentrations in the cell cytoplasm, causing toxicity that hampers plant growth [43]. Saffan noted that osmotic effects from salinity stress can disrupt the plant's water balance, leading to reduced turgor and inhibiting plant growth [44]. Similar reduction in agronomic characters of NaCl treated plants of rice varieties have been reported by many researchers [45,46,47].

Osmotic stress can result from high salinity, and high salt concentrations in soils make it more difficult for plants to absorb water and nutrients [48]. At the flowering stage, high salt makes pollens less viable, which affects grain yield [49,50].

In the present research work, we induced salt tolerance in salt sensitive varieties of Basmati rice using *in vitro* step up NaCl treatment technique and M<sub>2</sub> and M<sub>3</sub> generations of these Basmati varieties were harvested at 14 ds/m EC. According to IRRI, soil salinity beyond 4ds/m is considered as moderate salinity while more than 8 ds/m is high salinity for rice [12]. With increasing NaCl concentration, various agronomic and morphological characteristics of these salt-tolerant M<sub>2</sub> and M<sub>3</sub> generations significantly declined. Results of this research work led us to conclude that this induction of salt tolerance in these varieties of Basmati rice was due to the adaptation of calli to saline environment by step up salt treatment method.

## Acknowledgements

The present research work was funded by the Higher Education Commission (HEC), and all the laboratory facilities were provided by the Institute of Botany, University of the Punjab, Lahore, Pakistan.

## Author Contributions

Supervised, checked and proofreading by Dr. Humera Afrasiab.

Study design and conception, material preparation, data collection and statistical analysis, interpretation

of the results and first draft was written by Naveeda Batool.

## Conflict of Interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

## References

1. Joshi RK, Rao GJN. Somaclonal variation in submergence tolerant rice cultivars and induced diversity evaluation by PCR markers. *International Journal of Genetics and Molecular Biology*, (2009); 1(5): 80-88.
2. Shitu GA, Nain MS, Singh R. Developing extension model for uptake of Precision conservation agricultural practices in developing nations: learning from rice-wheat system of Africa and India. *Current Science*, (2018); 114(4): 814-825.
3. FAOSTAT: Food and Agriculture Organization of the United Nations. (November, 2022). Production/Yield Quantities of Rice, paddy in World. <http://www.fao.org/faostat/en/#data/QC/visualize>
4. Shahbandeh M. World rice acreage 2010-2018. (2020). <https://www.statista.com/statistics/271969/world-rice-acreage-since-2008/>
5. Chauhan BS, Jabran K, Mahajan G. (2017). Rice Production Worldwide. (2017); Springer International Publishing AG, Switzerland.
6. Economic Survey of Pakistan 2019–2020. (2020). [http://www.finance.gov.pk/survey/chapter\\_20/02\\_Agriculture.pdf](http://www.finance.gov.pk/survey/chapter_20/02_Agriculture.pdf)
7. Mahmood MA, Sheikh AD, Kashif M. Acreage supply response of rice in Punjab, Pakistan. *Journal of Agricultural Research*, (2007); 45(3): 231-236.
8. Grover A, Minhas D. Towards production of abiotic stress tolerant transgenic rice plants: issues, progress and future research needs. In: *Proc: Indian National Science Academy. Part B. Reviews and Tracts. Biological Sciences*, (2000); 66 (1): 13-32.
9. Ansari MR, Shaheen T, Bukhari SA, Husnain, T. Genetic improvement of rice for biotic and abiotic stress tolerance. *Turkish Journal of Botany*, (2015); 39: 911-919.
10. Fogliatto S, Serra F, Patrucco L, Milan M, Vidotto F. Effect of different water salinity levels on the germination of Imazamox-resistant and sensitive weedy rice and cultivated rice. *Agronomy*, (2019); 9: 658-670.
11. IRRI. (1997). Screening Rice for Salinity Tolerance. G.B. Gregorio, D. Senadhira & R.D. Mendoza. IRRI Discussion Paper Series No. 22.
12. IRRI. (2006). Bringing Hope, Improving Lives: Strategic Plan 2007–2015. Manila.p 61.
13. Acosta-Motos JR, Ortuno M, Bernal-Vicente A, Diaz-Vivancos P, Sanchez-Blanco MJ, Hernandez JA. Plant responses to salt stress: adaptive mechanisms. *Agronomy*, (2017); 7: 18.
14. Khan MA, Abdullah Z. Salinity-sodicity induced changes in reproductive physiology of rice (*Oryza sativa*) under dense soil conditions. *Environmental and Experimental Botany*, (2003); 49: 145-157.
15. Cha-um S, Trakulyingcharoen T, Smitamana P, Kirdmanee C. Salt tolerance in two rice cultivars differing in salt tolerant abilities in responses to Iso-osmotic stress. *Australian Journal of Crop Science*, (2009); 3(4): 221-230.
16. Qureshi RH, Aslam M, Mustafa G, Akhtar J. Some aspects of physiology of salt tolerance in wheat (*Triticum aestivum* L). *Pakistan Journal of Agricultural Sciences*, (1991); 28: 199-209.
17. Aslam M, Ahmad I, Mahmood IA, Akhtar J, Nawaz S. Physiological basis of differential tolerance in rice to

- Salinity. Pakistan Journal of Soil Sciences, (1995); 10: 38-41.
18. Shereen A, Mumtaz S, Raza S, Khan MA, Solangi S. Salinity effects on seedling growth and yield components of different inbred rice lines. Pakistan Journal of Botany, (2005); 37(1): 131-139.
19. Agricultural Statistics of Pakistan. (2019-20). Government of Pakistan, Ministry of Food, Agriculture and Livestock, Economic Wing, Islamabad.
20. Samal P, Babu S. The Shape of Rice Agriculture towards 2050. International Association of Agricultural Economists (IAAE), (2018); Conference, July 28-August 2, 2018, Vancouver, British Columbia. <https://ageconsearch.umn.edu/record/277550>
21. Nikam AA, Devarumath RM, Ahuja A, Babu H, Shitole MG, Suprasanna P. Radiation-induced in vitro mutagenesis system for salt tolerance and other agronomic characters in sugarcane (*Saccharum officinarum* L.). The Crop Journal, (2015); 3: 46-56.
22. Miki Y, Hashiba M, Hisajima S. Establishment of salt stress tolerant rice plants through step up NaCl treatment in vitro. Biologia Plantarum, (2001a); 44(3): 391-395.
23. Miki Y, Katoh M, Hisajima S. Salt tolerance of in vitro established salt-tolerant rice plants during further growth in soil. Biologia Plantarum, (2001b); 44(3): 463-466.
24. Tam DM, Lang NT. In vitro selection for salt tolerance in rice. Omonrice, (2003); 11: 68-73.
25. Sharry SE, Silva JAT. Effective organogenesis, somatic embryogenesis, and salt tolerance induction in vitro in the Persian Lilac tree (*Melia azedarach* L.). Floriculture, Ornamental and Plant Biotechnology, (2006); 2: 317-324.
26. Mahmood A, Latif T, Khan MA. Effect of salinity on growth, yield and yield components in Basmati rice germplasm. Pakistan Journal of Botany, (2009); 41(6): 3035-3045.
27. Murashige T, Skoog F. A revised medium for rapid growth and bioassay with Tobacco tissue cultures. Physiologia Plantarum, (1962); 15: 473-497.
28. Gupta B, Huang B. Mechanism of salinity tolerance in plants: Physiological, biochemical and molecular characterization. International Journal of Genomics, (2014); Article ID 701596, 18.
29. Ashraf M. Some important physiological selection criteria for salt tolerance in plants. Flora, (2004); 199: 361-376.
30. Rai MK, Kalia RK, Singha R, Gangolaa MP, Dhawana, AK. Developing stress tolerant plants through in vitro selection—An overview of the recent progress. Environmental and Experimental Botany, (2010); 71: 89-98.
31. Ullah H, Ullah I, Jadoon SA, Rashid H. Tissue culture techniques for callus induction in rice. Sarhad Journal of Agriculture, (2007); 23(1): 81-86
32. Afrasiab H, Jafar R. Effect of different media and solidifying agents on callogenesis and plant regeneration from different explants of rice (*Oryza sativa* L.) varieties Super Basmati and IRRI-6. Pakistan Journal of Botany, (2011); 43(1): 487-501.
33. Rafique MZ, Rashid H, Chaudhary MF, Chaudhry Z, Cheema NM. Study on callogenesis and organogenesis in local cultivars of rice (*Oryza sativa* L.). Pakistan Journal of Botany, (2011); 43(1): 191-203.
34. Singh S, Kumar A, Rana V, Kumar A, Sharma VR. In Vitro callus induction and plant regeneration in basmati rice (*Oryza sativa* L.) varieties. Advances in Agriculture and Natural Sciences for Sustainable. Journal of Pharmacognosy and Phytochemistry, (2018); SP5: 65-69.
35. Mostafiz SB, Wagiran A. Efficient callus induction and regeneration in selected Indica rice. Agronomy, 2018; 8:77.
36. Revathi S, Pillai MA. In vitro callus induction in rice (*Oryza sativa* L.). Research in Plant Biology, (2011); 1: 13-15.
37. Sikder MBH, Sen PK, Mamun MA, Ali MR, Rahman SM. In Vitro Regeneration of Aromatic Rice (*Oryza sativa* L.). International Journal of Agricultural and Biology, (2006); 6: 759-762.
38. Ge XJ, Chu JH, Lin YJ, Wang SP. A tissue culture system for different germplasms of indica rice. Plant Cell Reports, (2006); 25: 392-402.
39. Upadhyaya G, Sen M, Roy A. In vitro callus induction and plant regeneration of rice (*Oryza sativa* L.) var. 'Sita', 'Rupali' and 'Swarna Masuri'. Asian Journal of Plant Science and Research, (2015); 5: 24-27.
40. Htwe NN, Maziah M, Ling HC, Zaman, FQ, Zain AM. Responses of some selected Malaysian rice genotypes to callus induction under in vitro salt stress. African Journal of Biotechnology, (2011); 10: 350-362.
41. Wu J, Seliskar DM, Gallagher JL. The response of plasma membrane lipid composition in callus of the halophyte *Spartina patens* (Poaceae) to salinity stress. American Journal of Botany, (2005); 92: 852-858.
42. Queiros F, Fidalgo F, Santos I, Salem R. In vitro selection of salt tolerant cell lines in *Solanum tuberosum* L. Biologia Plantarum, (2007); 51(4): 728-734.
43. Khorami A, Safarnejad A. In vitro selection of *Foeniculum vulgare* for salt tolerance. Notulae Scientia Biologicae, (2011); 3: 90-97.
44. Saffan SES. Effect of salinity and osmotic stresses on some economic plants. Research Journal of Agriculture and Biological Sciences, (2008); 4: 159-166.
45. R. Munns, Tester, M. Mechanisms of salinity tolerance. Annual Review of Plant Biology, (2008); 59: 651-681.
46. Hakim MA, Juraimi AS, Begum M, Hanafi MM, Ismail MR, Selamat A. Effect of salt stress on germination and early seedling growth of rice (*Oryza sativa* L.). African Journal of Biotechnology, (2010); 9(13): 1911-1918.
47. Hussain S, Cao X, Zhong C, Zhu L, Khaskheli MA, et al. Sodium chloride stress during early growth stages altered physiological and growth characteristics of rice. Chilean Journal of Agricultural Research, (2018); 78(2): 183-197.
48. Verslues PE, Agarwal M, Katiyar-Agarwal S, Zhu JH, Zhu, JK. Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. The Plant Journal, (2006); 45(4): 523-539.
49. Khatun S, Flowers, TJ. Effects of salinity on seed set in rice. Plant Cell Environment, (1995); 18(1): 61-67.
50. Singh, RK, Mishra B, Singh KN. Salt tolerant rice varieties and their role in reclamation programme in Uttar Pradesh. Indian Farming, (2004); 6-10.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License. To read the copy of this license please visit: <https://creativecommons.org/licenses/by-nc/4.0/>